

## Harvesting methods

The age of trees when leaves can be harvested for processing into leaf powder is variable and mainly depends essentially on site conditions. Trees can be harvested from any age. In general, leaf utilisation could start before the sixth year when site conditions are favourable. Women traditionally start harvesting when leaves begin to develop and the period varies according to agroecological zones (April to May). Mass leaf harvesting is done in September and October. The bark is also harvested at the same time as the leaves. The fruit is harvested when the shell is brown, between December and April. The tools used in harvesting the leaves are the sickle and dolé. Harvesting by hand picking is done less frequently since it is difficult to climb a baobab tree.

## Acknowledgement

Members of Agri-Africa/Karwil Consultancy are herewith acknowledged for the information provided.

## References

- Van Wyk, B-E., Van Oudtshoorn & Gericke, N. 1997. *Medicinal plants of South Africa*, Pretoria: Briza Publications.  
Hardy, D.S. 1984. Muvhuyu—the baobab; *Orbea verrucosa* var *fucosa*. *Aloe*, 21: 43–45. The National Library of South Africa.  
<http://www.baobabfruitco.com>  
[http://www.icuc-iwmi.org/files/Publications/Baobab\\_manual.pdf](http://www.icuc-iwmi.org/files/Publications/Baobab_manual.pdf)

## Uses

Baobab provides food, emergency water and fibre. It has also medicinal uses. Fibre from the stringy inner bark provides items (or is used for items) such as rope, thread, basket, nets, snares, fishing lines, strings for musical instruments, and a paper stock tough enough for banknotes. The fibre is even used for weaving. Some is woven into fabrics that are valued for making the bags used for carrying and storing everyday goods. Baobab trees supply food and traditional medicines for both humans and their livestock. A refreshing drink, prepared from the pale yellow or whitish fruit pulp called cream of tartar, has been used to treat fevers, diarrhea and apparently also haemoptysis. The leaves are used against fever, to reduce perspiration and as an astringent. They also come in handy in treating other afflictions: asthma, kidney and bladder diseases, insect bites, fevers, malaria and sores.

In the Limpopo Province the powdered seeds are given to children as a hiccup remedy. Under survival stress man can use many parts of the baobab as food or obtain water from its roots, branches or leaves. A crude, coffee-like beverage can be prepared by baking the baobab seeds. Many people find shelter from the blistering sun in the ample shade provided by its sturdy trunk. Young leaves when mixed with pepper and salt and added to a stew give it a good taste. Young, fresh leaves are cut into pieces and cooked into a sauce. Sometimes the leaves are dried and powdered and used for cooking.

# Baobab



Further information can be obtained from

Directorate Plant Production  
Private Bag X250  
PRETORIA 0001  
Tel +27 12 319 6072  
Fax +27 12 319 6079  
Email DPPP@daff.gov.za

2012

Printed and published by

Department of Agriculture, Forestry and Fisheries

Obtainable from

Resource Centre  
Directorate Communication Services  
Private Bag X144  
PRETORIA  
0001

agriculture,  
forestry & fisheries

Department:  
Agriculture, Forestry and Fisheries  
REPUBLIC OF SOUTH AFRICA



## Planting

picking by overzealous individuals (for soup leaves) but mature trees have few enemies. Neither cattle nor goats do serious harm. Not even overzealous pickers can seemingly set back a healthy old baobab. There are no serious pests and diseases of baobab. However, some fungal and viral diseases have been recorded and several insects attack the wood, fruit and young shoots. The most investigated common pests are:

- cotton bollworms *Heliothis armigera*, *Diparopsis castanea* and *Earias biplaga*;
- cotton-stainers (bugs) such as *Dysdercus fasciatus*, *D. intermedius*, *D. nigrofasciatus*, *D. superstitiosus*, *Odontotarsus exsanguinis*, *O. sexpunctatus*;
- *Oxycarenus albipennis* as well as flea beetles, *Podagrica* spp.
- *Symptoms*  
The newly emerged larvae feed on the leaf and foliage of the plants. Cotton bollworms tunnel into the fruits of the baobab. They suck the sap of the leaves and young foliage. The immature fruit drops.

### CONTROL

Registered chemical fertilisers are recommended for use. A decoction of the kernels of *Azadirachta indica* (neem) can be used for insect control. Weeding can also be used as a control measure.

The baobab is a host for members of the Pseudococcidae, the mealybugs, which can be vectors for virus diseases of cocoa and the cocoa capsid, *Diastictica theobromae*. In the Limpopo and Mpumalanga provinces of South Africa, a caterpillar, *Gonimbrasia herlina*, can feed on the leaves.

### SYMPTOMS

The caterpillar sucks the sap of the leaflets, mature and tender shoots, leaf petiole bases and young foliage. The immature fruit drops. Chlorotic leaves and defoliation can be observed. The caterpillar feeds on the leaves.

### CONTROL

The affected parts should be removed. The caterpillars can be removed by hand and crushed.

## Scientific name: *Adansonia digitata*

Common names: Baobab, Monkey bread, Ethiopian sour gourd, Cream-of-tartar tree (English); Kremetartboom (Afrikaans); Muvhuyu (Tshivenda); Shimuwu (isiTsonga); Isimuhu, Umshimulu (isiZulu); Mowana (Setswana); Moyo (Northern Sotho).

## Background

Baobab is a deciduous tropical fruit tree ranging in height from 5 to 25 m and is distributed in belts in low-lying areas across Africa, Madagascar, India, Sri Lanka and Australia. It belongs to the family called Bombacaceae. It is a very long-lived, fast-growing tree and has a lifespan of hundreds to thousands of years. The growth of the baobab is mainly managed and protected by local people. Baobabs are widespread throughout the hot, drier regions of tropical Africa, extending from Mozambique, the northern provinces of South Africa and Namibia to Ethiopia, Sudan and southern fringes of the Sahara. In South Africa the tree is found in the frost-free areas near Waterpoort in the Western Cape and the Limpopo Province.

## Climatic and soil requirements

Baobabs occur in semiarid to subhumid tropical zones. They grow on many different soils including sandy loam but develop best on calcareous substrates and on deep, slightly moist sites. They thrive where the average annual temperature is 20 to 30 °C. Germination is achieved only when soil temperature exceeds 28 °C. Baobabs are extremely susceptible to frost throughout their life cycle.

## Cultural practices

### Soil preparation

Land preparation is done in the summer or at the onset of the rainy season to preserve the soil structure. The soil should be ploughed 3 to 4 weeks prior to transplanting, then again after 15 days, and then again just before planting the seedlings. The soil should be leveled and have good drainage.

Seedlings are mainly raised and transplanted into the field at 10 m x 10 m spacing. The hole size is 60 cm x 60 cm x 60 cm, but smaller may be suitable (40 cm<sup>3</sup>). The trees, planted in a row, should be given weekly volume of water which vary from 10 ℥ for the first tree, 15 ℥ for the second and 30 ℥ for the third. Planting is done when the rainy season has started. Cuttings should be 5 cm to 10 cm in length and pushed straight into the soil to a depth of about 2.5 cm. For leaf production only, planting should be done 0.2 m x 0.5 m and for leaves and fruit it should be 4 m x 4 m.

## Propagation

Baobabs can be propagated from seeds as well as vegetatively. Vegetative propagation involves the growth of a new tree from a shoot, bud or cutting from a good-quality mature tree. The trees have traditionally been propagated by transplanting naturally regenerated seedlings.

## Fertilisation

Organic and mineral fertilisers can be used and it is recommended to use farmyard manure, compost or green legume manures, especially at the time of planting for intensive leaf production.

### Irrigation

The volume of water required varies with the size of the tree and is dependent on local climate. In general, during establishment, about 1 ℥ to 2 ℥ of water should be applied twice a week to the base of each young tree. The small trees can be irrigated regularly to produce higher returns from intensive leaf production and better growth. Mature baobab trees require no irrigation.

## Weed control

The weeds should be removed from around the tree during the early stages of growth.

## Disease and pest control

Few small baobabs are ever seen nowadays because they fall victim to grazing by cattle and goats, ground fires, or